

Real divergence-free Jacobian maps are shear maps *

M. Sabatini

May 18, 2009

Abstract

We show that divergence-free jacobian maps, recently considered in [11], [14], are actually shear maps.

1 Introduction

Let $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a continuously differentiable map. By the inverse function theorem, if its Jacobian determinant $\det J_\Phi$ does not vanish at a point, then Φ is locally invertible at such a point. It is well-known that the local invertibility of Φ , even if it holds on all of $\mathbb{R}^n$, does not imply Φ 's global invertibility. Additional conditions implying Φ 's global invertibility, or just injectivity, have been studied in several fields (see [1], [13]). Some conditions have just been conjectured to be sufficient to guarantee the map's invertibility, as in the case of the celebrated Jacobian Conjecture [8]. Originally formulated for complex maps, and successively studied for maps defined on arbitrary fields, such a conjecture asks to prove that a polynomial map with constant non-zero Jacobian determinant is globally invertible, with polynomial inverse. The problem has been attacked in several ways. One can find in [2] an overview of results concerning the Jacobian Conjecture up to 1982. A more recent account is contained in [3]. Among general results concerning such a problem, it is known that it is equivalent to prove or disprove the statement in any field of zero characteristic, that it is sufficient to prove Φ 's injectivity in order to get its surjectivity [2], and that Φ 's global invertibility implies that Φ^{-1} is a polynomial map. The most studied special case is the bidimensional one, $\Phi(x, y) = (P(x, y), Q(x, y))$, where the statement was proved under the hypothesis that either P 's or Q 's degree is

*2000 AMS *Mathematics Subject Classification*: 14R15. *Key words and phrases*: Jacobian Conjecture, divergence-free, shear map.

4, or prime, or both degrees are ≤ 100 (see [2] for a more comprehensive list of results).

Recently, a particular class of maps has been taken into account, namely the so-called divergence-free maps [11]. Such maps have the form $\Phi(x, y) = (x + p(x, y), y + q(x, y))$, where $p(x, y)$ and $q(x, y)$ are polynomials of order > 1 , with $\frac{\partial p}{\partial x} + \frac{\partial q}{\partial y} = 0$. For such maps one has $\det J_\Phi = 1 + \left(\frac{\partial p}{\partial x} + \frac{\partial q}{\partial y}\right) + \left(\frac{\partial p}{\partial x} \frac{\partial q}{\partial y} - \frac{\partial p}{\partial y} \frac{\partial q}{\partial x}\right)$, hence under the condition $\frac{\partial p}{\partial x} + \frac{\partial q}{\partial y} = 0$ also the determinant-like term vanishes, $\frac{\partial p}{\partial x} \frac{\partial q}{\partial y} - \frac{\partial p}{\partial y} \frac{\partial q}{\partial x} = 0$. In algebraic terms, this amounts to ask that no terms in the divergence-like part of $\det J_\Phi$ cancels with any term in its determinant-like part (maps with a linear part different from the identity can be reduced to the above form by multiplying by an invertible linear map). Such a condition is satisfied under some symmetry conditions on $p(x, y)$ and $q(x, y)$, as in the case of $p(x, y)$ and $q(x, y)$ even polynomials, since the terms in $\frac{\partial p}{\partial x} + \frac{\partial q}{\partial y}$ have odd degree, while the terms in $\frac{\partial p}{\partial x} \frac{\partial q}{\partial y} - \frac{\partial p}{\partial y} \frac{\partial q}{\partial x}$ have even degree. In [11] it was proved that such maps are globally invertible. A different proof for real maps, with some extensions, has been given in [14], by applying results proved in relation to the Global Asymptotic Stability Jacobian Conjecture [5], [6], [7].

In this paper we give a complete characterization of real divergence-free maps, proving that every such map has the form

$$\Phi(x, y) = \left(x + \sum_{i=2}^n \varepsilon_i \alpha (\beta x - \alpha y)^i, y + \sum_{i=2}^n \varepsilon_i \beta (\beta x - \alpha y)^i \right), \quad (1)$$

with $\alpha, \beta \in \mathbb{R}$, $\varepsilon_i \in \mathbb{R}$ for $i = 2, \dots, n$. This means that the nonlinear part of such maps essentially depends on the single variable $\beta x - \alpha y$, so that after a change of variables the map has the form of a shear map, $(u, v) \mapsto (u, v - f(u))$. Our approach consists in solving the system of PDEs

$$\frac{\partial p}{\partial x} + \frac{\partial q}{\partial y} = 0, \quad \frac{\partial p}{\partial x} \frac{\partial q}{\partial y} - \frac{\partial p}{\partial y} \frac{\partial q}{\partial x} = 0, \quad (2)$$

in the class of homogeneous polynomials, then proving that only polynomials with nonlinear part as in (1) solve the system (2). As a consequence, we prove that under some simple algebraic conditions, a jacobian map is the composition of a linear automorphism and a shear map.

2 Results

Let $\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $\Phi(x, y) = (P(x, y), Q(x, y))$ be a real polynomial map. Let J_Φ be its jacobian matrix. We say that Φ is a *jacobian map* if its jacobian determinant $\det J_\Phi$ is a non-zero constant. Possibly replacing $\Phi(x, y)$ with

$\Phi(x, y) - \Phi(0, 0)$, which has the same jacobian determinant, we may assume that $\Phi(0, 0) = (0, 0)$. Moreover one can compose Φ with the inverse matrix of $J_\Phi(0, 0)$, obtaining another jacobian map with linear part coinciding with the identity. Let us call again Φ the map obtained by such operations. Then we can write

$$P(x, y) = x + \sum_{i=2}^n p_i(x, y), \quad Q(x, y) = y + \sum_{i=2}^n q_i(x, y),$$

where $p_i(x, y)$, $q_i(x, y)$ are homogeneous polynomials of degree i . Let us set $\bar{\Phi}(x, y) = \Phi(x, y) - (x, y)$. In other words, $\bar{\Phi}$ is the map consisting only of the nonlinear terms of Φ . We say that $\Phi(x, y)$ is a *divergence-free jacobian map* if it satisfies the condition

$$\frac{\partial}{\partial x} \sum_{i=2}^n p_i(x, y) + \frac{\partial}{\partial y} \sum_{i=2}^n q_i(x, y) = 0.$$

As observed in the introduction, if Φ satisfies the above condition, then $\det J_\Phi = 0$.

We say that Φ is a *shear map* if its nonlinear terms are linear combinations of powers of a homogeneous polynomial of degree 1, as for the map $\Phi(x, y) = (3x - 4y + (x - y)^2, -2x + y + (x - y)^2)$. The adjective *shear* comes from the fact the nonlinear part of that such maps substantially acts as a displacement along a given direction. The displacement's amount depends nonlinearly on the position of (x, y) .

In next lemma we show that a homogeneous polynomial satisfying simultaneously the divergence and the determinant condition is essentially a function of a single variable. We start considering homogeneous polynomials.

Lemma 1 *Let $p(x, y)$, $q(x, y)$ be homogeneous polynomials of the same degree $j > 1$, not both identically zero. If*

$$\frac{\partial p}{\partial x} + \frac{\partial q}{\partial y} = 0, \quad \frac{\partial p}{\partial x} \frac{\partial q}{\partial y} - \frac{\partial p}{\partial y} \frac{\partial q}{\partial x} = 0, \quad (3)$$

then there exist $\alpha, \beta, \zeta \in \mathbb{R}$, $\alpha^2 + \beta^2 \neq 0$, $\zeta \neq 0$, such that $p(x, y) = \zeta \alpha (\beta x - \alpha y)^j$, $q(x, y) = \zeta \beta (\beta x - \alpha y)^j$.

Proof. Let us assume $q(x, y)$ not to be constant. Then there exists a line $l = \{(x, y) : x = t \cos \theta, y = t \sin \theta\}$ such that $q(t \cos \theta, t \sin \theta)$ is non-constant. The gradient ∇q does not vanish at any point of l . By the second equality in (3), in a neighbourhood of the point $(\cos \theta, \sin \theta)$ there exists an analytic function ψ such that $p(x, y) = \psi(q(x, y))$.

Then there exist $\alpha, \beta \in \mathbb{R}$, $\alpha^2 + \beta^2 \neq 0$ such that

$$\alpha t^j = p(t \cos \theta, t \sin \theta) = \psi(q(t \cos \theta, t \sin \theta)) = \psi(\beta t^j).$$

One has $\beta \neq 0$, otherwise $q(t \cos \theta, t \sin \theta) = \beta t^j$ would be constant. Setting $s = \beta t^j$, one has $t^j = \frac{s}{\beta}$, hence

$$\psi(s) = \psi(\beta t^j) = \alpha t^j = \frac{\alpha s}{\beta}.$$

This proves that locally $\beta p(x, y) - \alpha q(x, y) = 0$. By the identity principle for polynomials, the same equality holds on all of $\mathbb{R}^2$. As a consequence, there exist a homogeneous polynomial $r(x, y)$ of degree j and $k \in \mathbb{R}$, $k \neq 0$, such that $p(x, y) = k\alpha r(x, y)$, $q(x, y) = k\beta r(x, y)$. Then one has

$$0 = \frac{\partial p}{\partial x} + \frac{\partial q}{\partial y} = k\alpha \frac{\partial r}{\partial x} + k\beta \frac{\partial r}{\partial y} = k(\alpha, \beta) \cdot \nabla r(x, y).$$

This implies that the level curves of $r(x, y)$ are just the lines $\beta x - \alpha y = \text{const.}$, hence there exists a function ρ such that $r(x, y) = \rho(\beta x - \alpha y)$. Choosing $(x, y) = \left(\frac{\beta t}{\alpha^2 + \beta^2}, -\frac{\alpha t}{\alpha^2 + \beta^2}\right)$ one has $\beta x - \alpha y = t$ and

$$\rho(t) = r\left(\frac{\beta t}{\alpha^2 + \beta^2}, -\frac{\alpha t}{\alpha^2 + \beta^2}\right) = ht^j.$$

for some $h \in \mathbb{R}$, $h \neq 0$. In conclusion, $r(x, y) = \rho(\beta x - \alpha y) = h(\beta x - \alpha y)^j$. Setting $\zeta = hk$ gives the statement. $\clubsuit$

Remark 1 *The above lemma shows that a divergence-free jacobian map can be written as follows,*

$$P(x, y) = x + \sum_{i=2}^n \zeta_i \alpha_i (\beta_i x - \alpha_i y)^i, \quad Q(x, y) = y + \sum_{i=2}^n \zeta_i \beta_i (\beta_i x - \alpha_i y)^i, \quad (4)$$

with $\zeta_i \neq 0$, $\alpha_i^2 + \beta_i^2 \neq 0$ for $i = 0, \dots, n$. In fact, since “divergence” terms do not cancel with “determinant” terms, every homogeneous couple $(p_j(x, y), q_j(x, y))$ satisfies the hypotheses of lemma 1.

Something more can be said comparing terms with different degrees. Next lemma is concerned with couples of arbitrary functions both depending only on a first-degree homogeneous polynomial.

Lemma 2 *Let us set $p(x, y) = a\phi(\beta x - \alpha y) + b\psi(\delta x - \gamma y)$, $q(x, y) = c\phi(\beta x - \alpha y) + d\psi(\delta x - \gamma y)$, with ϕ, ψ non-constant functions of class C^1 . If (3) holds, then the lines $\beta x - \alpha y = 0$ and $\delta x - \gamma y = 0$ coincide.*

Proof. One has

$$0 = \frac{\partial p}{\partial x} \frac{\partial q}{\partial y} - \frac{\partial p}{\partial y} \frac{\partial q}{\partial x} = (ad - bc)(\alpha\delta - \beta\gamma)\phi'(\beta x - \alpha y)\psi'(\delta x - \gamma y).$$

At every point where both ϕ' and ψ' do not vanish one has $(ad-bc)(\alpha\delta-\beta\gamma) = 0$. If $\alpha\delta-\beta\gamma = 0$, then the thesis is proved. If $ad-bc = 0$, then there exist $m, \eta, \omega \in \mathbb{R}$, $m \neq 0$, $\eta^2 + \omega^2 \neq 0$ and a function $r(x, y)$ such that $p(x, y) = m\eta r(x, y)$, $q(x, y) = m\omega r(x, y)$. Then, as in lemma 1, one has

$$0 = \frac{\partial p}{\partial x} + \frac{\partial q}{\partial y} = m\eta \frac{\partial r}{\partial x} + m\omega \frac{\partial r}{\partial y} = m(\eta, \omega) \cdot \nabla r(x, y).$$

As in lemma 1, this proves that there exists one-variable function $\rho(t)$ such that $r(x, y) = \rho(\omega x - \eta y)$. Now, assume by absurd that the lines $\beta x - \alpha y = 0$ and $\delta x - \gamma y = 0$ do not coincide. Then one can make an invertible change of variables, $u = \beta x - \alpha y$, $v = \delta x - \gamma y$, and consider the functions of u and v obtained from $p(x, y), q(x, y), r(x, y)$ by applying the change of variables,

$$p^*(u, v) = a\phi(u) + b\psi(v) = m\eta\rho(k_1u + k_2v),$$

$$q^*(u, v) = c\phi(u) + d\psi(v) = m\omega\rho(k_1u + k_2v),$$

where $k_1, k_2 \in \mathbb{R}$, $k_1^2 + k_2^2 \neq 0$. Differentiating both equalities with respect to u one has

$$a\phi'(u) = m\eta k_1 \rho'(k_1u + k_2v), \quad c\phi'(u) = m\omega k_1 \rho'(k_1u + k_2v).$$

If $k_2 \neq 0$, then the above equalities hold only if ϕ' and ρ' are constant, against the hypothesis, hence $k_2 = 0$. Similarly, differentiating with respect to v one proves that $k_1 = 0$. This leads to $k_1 = 0 = k_2$, contradiction. ♣

Theorem 1 *Let $\Phi(x, y) = (P(x, y), Q(x, y))$ be a divergence-free jacobian map of degree n . Then there exist $\alpha, \beta \in \mathbb{R}$, $\alpha^2 + \beta^2 \neq 0$, $\varepsilon_i \in \mathbb{R}$, such that*

$$P(x, y) = x + \sum_{i=2}^n \varepsilon_i \alpha (\beta x - \alpha y)^i, \quad Q(x, y) = y + \sum_{i=2}^n \varepsilon_i \beta (\beta x - \alpha y)^i, \quad (5)$$

Proof. Let n be the degree of Φ . Due to lemma 1, $P(x, y)$ and $Q(x, y)$ are as in remark 1. Let us set $\alpha = \alpha_n$, $\beta = \beta_n$, $\varepsilon_n = \zeta_n$. Since we assume Φ to be of degree n , $\alpha^2 + \beta^2 \neq 0$, $\varepsilon_n \neq 0$.

We claim that for $i = 2, \dots, n-1$, $\alpha_i = \alpha$, $\beta_i = \beta$.

Let us proceed by decreasing degree. The functions $p(x, y) = \zeta_n \alpha (\beta x - \alpha y)^n + \zeta_{n-1} \alpha_{n-1} (\beta_{n-1} x - \alpha_{n-1} y)^{n-1}$, $q(x, y) = \zeta_n \beta (\beta x - \alpha y)^n + \zeta_{n-1} \beta_{n-1} (\beta_{n-1} x - \alpha_{n-1} y)^{n-1}$ satisfy the hypotheses of lemma 2, since their terms do not interact with those ones of lower degrees. Then, by lemma 2, the terms $\beta x - \alpha y$ and $\beta_{n-1} x - \alpha_{n-1} y$ are proportional. Since $\beta x - \alpha y$ does not vanish identically, there exist $\xi_{n-1} \in \mathbb{R}$ such that $\alpha_{n-1} = \xi_{n-1} \alpha$, $\beta_{n-1} = \xi_{n-1} \beta$. As a consequence, the terms of degree $\geq n-1$ in $P(x, y)$ and $Q(x, y)$ are all function of $\beta x - \alpha y$. Setting $\varepsilon_{n-1} = \zeta_{n-1} \xi_{n-1}^n$, the form of the $(n-1)$ -degree term in (4) is as in (5).

Now we may repeat the procedure replacing $n-1$ with $n-2$. We apply again lemma 2 to the functions $p(x, y) = \alpha \left(\varepsilon_n (\beta x - \alpha y)^n + \varepsilon_{n-1} (\beta x - \alpha y)^{n-1} \right) + \zeta_{n-2} \alpha_{n-2} (\beta_{n-2} x - \alpha_{n-2} y)^{n-2}$, $q(x, y) = \beta \left(\varepsilon_n (\beta x - \alpha y)^n + \varepsilon_{n-1} (\beta x - \alpha y)^{n-1} \right) + \zeta_{n-2} \beta_{n-2} (\beta_{n-2} x - \alpha_{n-2} y)^{n-2}$, in order to prove that $\beta x - \alpha y$ and $\beta_{n-2} x - \alpha_{n-2} y$ are proportional. Since $\beta x - \alpha y$ does not vanish identically, there exist $\xi_{n-2} \in \mathbb{R}$ such that $\alpha_{n-2} = \xi_{n-2} \alpha$, $\beta_{n-2} = \xi_{n-2} \beta$.

Such a procedure can be applied $n-2$ times in order to prove that every nonlinear term in Φ is actually a power of $\beta x - \alpha y$ multiplied by a constant $\varepsilon_i = \zeta_i \xi_i^{i+1}$, $i \neq n$. $\clubsuit$

In the above theorem it is not assumed that for $i = 2, \dots, n-1$ one has $\varepsilon_i \neq 0$. Actually some degrees may not appear in a divergence-free jacobian map, as in $\Phi(x, y) = (x - y^2 - y^5, y)$.

In [11] and [14] some simple algebraic conditions ensuring a jacobian map to be divergence-free were taken into account. We apply them to the present situation.

Given a polynomial P , we write $d(P)$ for its degree, $o(P)$ for its order. We say that a polynomial is *even* if it is the sum of even degree monomials, *odd* if it is the sum of odd degree monomials. Similarly, we say that a polynomial is *x-even* if it contains only terms with even powers of x , *x-odd* if it contains only terms with odd powers of x . We say that a non-negative integer is a *gap* of P if it is the difference of the degrees of two distinct monomials in P . We denote by $G(P)$ the gap-set of P . As an example, the polynomial $P(x, y) = x^3 + y^3 + x^2 y^2 + y^7$ has gap-set $G(P) = \{0, 1, 3, 4\}$. If P has exactly one monomial, or if P is identically zero, then we say that it has empty gap-set.

We say that the couple of polynomials (P, Q) satisfies the *gap condition* if for every monomial M in P , one has $d(M) - 1 \notin G(Q)$. The gap condition is not symmetric, as shown by the couple $(P, Q) = (x + y^2, x^6 + y^2)$. In such a case one has $G(P) = \{1\}$, $G(Q) = \{4\}$, so that (P, Q) satisfies the gap condition, but (Q, P) does not.

We say that (P, Q) satisfies the *symmetric gap condition* if both (P, Q) and (Q, P) satisfy the gap condition.

Corollary 1 *Let $\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a jacobian map of the type $\Phi(x, y) = (ax + by + p(x, y), cx + dy + q(x, y))$, $a, b, c, d \in \mathbb{R}$, $o(p) > 1$, $o(q) > 1$. If one of the following holds,*

- i) $\max\{d(p), d(q)\} < o(p) + o(q) - 1$,*
- ii) both $p(x, y)$ and $q(x, y)$ are even polynomials,*
- iii) p is odd, q is even and (p, q) satisfies the gap condition,*

*iv) (p, q) satisfies the symmetric gap condition,
then $\Psi = \Phi \circ (J_\Phi(0, 0))^{-1}$ is a shear map.*

Proof. Let us consider the map $\Psi = \Phi \circ (J_\Phi(0, 0))^{-1}$. One has $\Psi(x, y) = (x + p^*(x, y), y + q^*(x, y))$, with $o(p^*) > 1$, $o(q^*) > 1$. The composition with a linear map does not change the properties in *i), ..., iv)*, hence Ψ satisfies all of them. In [14] it was proved that any of the above conditions implies Ψ to be a divergence-free map. Then the conclusion comes from theorem 1. ♣

Next corollary is concerned with a kind of symmetry which is not preserved by composition with linear transformations.

Corollary 2 *Let $\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a jacobian map of the type $\Phi(x, y) = (x + p(x, y), y + q(x, y))$, $o(p) > 1$, $o(q) > 1$. If one of the following holds,
i) p is x -even, q is x -odd,
ii) p is y -odd, q is y -even,
then Φ is a shear map.*

Proof. By theorem 2 of [14], Φ is a divergence-free jacobian map. Then one can apply theorem 1. ♣

References

- [1] A. Ambrosetti, G. Prodi, *A primer of nonlinear analysis. Corrected reprint of the 1993 original*, Cambridge Studies in Advanced Mathematics, **34**. Cambridge University Press, Cambridge, 1995.
- [2] H. Bass, E. H. Connell, D. Wright, *The jacobian conjecture: reduction of degree and formal expansion of the inverse*, Bull. Amer. Math. Soc. **7** (1982), 287–330.
- [3] M. de Bondt, A. van den Essen, *Recent progress on the Jacobian Conjecture*, Bull. Amer. Math. Soc. **7** (1982), 287–330.
- [4] L. Druzkowski, *An effective approach to Keller's Jacobian conjecture*, Math. Ann. **264**, **3** (1983), 303–313.
- [5] R. Fessler, *A proof of the two-dimensional Markus-Yamabe stability conjecture and a generalization*, Ann. Pol. Math. **62**, **1** (1995), 45–74.
- [6] A. A. Glutsyuk, *Complete solution of the Jacobian problem for planar vector fields (Russian)*, Uspekhi Mat. Nauk **49**, **3** (1994), 179–180, translation in Russian Math. Surveys **49**, **3** (1994) 185–186.

- [7] C. Gutierrez, *A solution to the bidimensional global asymptotic stability conjecture*, Ann. Inst. H. Poincaré Anal. Non Linéaire **12**, **6** (1995), 627–671.
- [8] O. H. Keller, *Ganze Cremona-Transformationen*, Monats. Math. Physik. **47** (1939), 299–306.
- [9] G. Meisters, *Jacobian problems in differential equations and algebraic geometry*, Rocky Mountain J. Math. **12** (1982), 679–705.
- [10] L. Markus, H. Yamabe, *Global stability criteria for differential systems*, Osaka Math. Jour. **12** (1960), 305–317.
- [11] J. W. Neuberger, *The divergence-free Jacobian conjecture in dimension two*, Rocky Mountain J. Math. **36** (2006), 265–271.
- [12] C. Olech, *On the global stability of an autonomous system in the plane*, Contr. to Diff. Equations **1** (1963), 389–400.
- [13] T. Parthasarathy, *On global univalence theorems*, Lecture Notes in Mathematics, **977**. Springer-Verlag, Berlin-New York, 1983.
- [14] M. Sabatini, *A note on the divergence-free Jacobian Conjecture in $\mathbb{R}^2$* , arXiv:0903.4626v1 [math.AG].